

remescar

stretch marks

Description

Stretch marks or striae are a form of scarring caused by tearing in the dermal layer of the skin. The most common sites to be affected by stretch marks include the abdomen, breasts, thighs, or buttocks. Stretch marks are often the result of the rapid stretching of the skin associated with weight gain (e.g. pregnancy, obesity) or rapid growth (e.g. puberty) or sickness (e.g. Cushing's disease).

Remescar stretch marks® is a silicone based cream which prevents and treats stretch marks by releasing a long lasting protective film that will cover the affected skin. This thin and discrete barrier has several active properties.

The product will protect the skin and moderate inflammation. It will lead to an excellent integration of the cream into the skin lipid barrier, thanks to the phospholipid-induced spherical film and hereby it will manage the micro-environment of the scar in order to stimulate the healing.

In addition, the moisturizing and reconstructive ingredients will generate immediate and long term hydration of the skin, and optimize the skin's natural moisture balance.

Thanks to these properties, Remescar stretch marks® will also reinforce and visually improve the skin surface.

Intended use

Remescar stretch marks® is designed for the prevention, management and treatment of old and new stretch marks. Remescar stretch marks® releases a transparent, thin and silicone based film which creates a barrier over the affected skin.

Instructions for use

The skin should be clean and dry. At first time use, open the tube and remove the protective cap. Remescar stretch marks® should be applied gently on the stretch marks or striae. Massage the cream in until completely absorbed. Do not rub too hard as this will not increase the effectiveness. Apply Remescar stretch marks® twice a day for 2 to 3 months.

As a preventive measure: use from the second month of pregnancy.

On existing stretch marks: use for 3 months continuously on a normal healthy skin.

Precautions

- For external use only
- Do not use the product in the following circumstances: on third degree burns, on a caesarean, on open or infected wounds, on a skin currently affected by acne.

- Avoid contact with mucous membranes or the eyes.
- Not to be used on dermatological conditions that disrupts the integrity of the skin.
- Remescar stretch marks® is not sterile and must not be used as a wound dressing.
- Use of the product is inadvisable if you have allergic reactions to one of the ingredients. If skin sensitization occurs, discontinue use and consult your physician if the skin sensitization persists.
- Not recommended for children under the age of 6 without supervision.
- If the product is being used on the breasts while breastfeeding it's advised to wait 30 minutes after applying the product and respect the basic hygiene before every feeding.
- It's not advised to apply Remescar stretch marks® over other skin treatments.
- Do not use after the expiry date. Once opened, it's advised to use the product not longer than 6 months.
- May stain clothes if not completely dry.
- It's advisable to store the product below 30°C.
- Keep out of the reach of children.

Ingredients

Aqua; Caprylic/Capric Triglyceride; Glyceryl Stearate, Behenyl Alcohol, Palmitic Acid, Stearic Acid, Lecithin, Lauryl Alcohol, Myristyl Alcohol, Cetyl Alcohol; Methylpropanediol; Xylityl Glucoside, Anhydroxylitol, Xylitol, Aqua; Isodecyl Neopentanoate; Glycerin; Coconut Alkanes, Cococaprylate/caprate; Betaine; PVP; Olea Europaea (olive) Oil Unsaponifiables, Tocopherol; Squalane; Phenoxyethanol, Ethylhexylglycerin; Chlorphenesin; Dimethicone; Butyrospermum Parkii; Carbomer; Aminomethyl Propanol; Xanthan Gum; Hydroxyethylcellulose; Hydrogenated Lecithin; Disodium EDTA

Symbols

-  = batch number
-  = expiry date
-  = read the instructions for use
-  = keep away from sunlight


Sylphar NV
Xavier De Cocklaan 42
9831 Deurle
Belgium
www.remescar.com


0197